LIFTMODE

LIFTMODE 47 W. Polk St. STE 100-241 Chicago, IL 60605 liftmode@liftmode.com www.liftmode.com

CERTIFICATE OF ANALYSIS

Dihydromyricetin

(Ampelopsin)

Material Lot #: 190604 Country of Origin: China Manufacturer Analysis Date: 06/04/2019

Analysis Date: 02/10/2020 Retesting Date: 02/10/2023

Analysis	Claim	Result
Dihydromyricetin	98%	100.6%
Test	Specification	Result
H-NMR ID HPLC Assay Arsenic Lead Cadmium Mercury Total Aerobic Count Yeast & Mold E.coli Salmonella Coliform	Conforms ≥98% ≤1.5 ppm ≤0.5 ppm ≤0.5 ppm ≤3 ppm <1,000 cfu/g <100 cfu/g Negative Negative <10 cfu/g	Conforms 100.6% 1.364 ppm 0.101 ppm <0.0001 ppm 0.005 ppm Conforms Conforms Conforms Conforms Conforms

Dihydromyricetin should be stored at or below room temperature in a tightly sealed durable container. Dihydromyricetin should be protected from excess heat, direct sunlight, excess humidity, and moisture. Dihydromyricetin has a retesting period of 3 years from the date of analysis when properly stored.

DHM

Dihydromyricetin

Main Benefits

- DHM has been found to be a powerful anti-inflammatory in its ability to regulate genes relating to inflammation and inhibit inflammation in HeLa cells
- Studies indicate that DHM may also show promise in promoting healthy levels of apoptosis.
- DHM has also been found to effectively reduce alcohol related hangovers and promote a healthy metabolism while simultaneously holding effective neuroprotective properties.

Main Cautions

- DHM is generally recognized as safe when taken as a dietary supplement. Rat studies with doses of up to 22g/1kg body weight failed to find any side-effects.
- Using the HED ratio, the safe upper limit dosage for DHM would be 15.68g for a 70kg person. However we strongly advise that you do not exceed the recommended serving size.
- Do not use this supplement without first consulting with your doctor if you are taking any medication or have any medical condition.

Usage Tips

- A 0.625 cc measuring scoop is included. One rounded scoop contains approximately one serving, or approximately 400mg of DHM. As a dietary supplement, take 1 serving 1-2 times per day. Start at the lower suggested quantity to assess response.
- The negative effects of DHM are dependent on the amount taken. Use of a scale with 10mg/0.01g accuracy or better is highly recommended.
- Use of capsules, or mixing with tea, yogurt, apple sauce, or oatmeal may help make the powder easier to tolerate.
- This supplement is not intended to treat, diagnose, prevent, or cure any diseases. Consult your healthcare provider before use if you have a medical condition or if you are taking any prescription medications.
- It is safe to stack DHM with other supplements, so long as the amount consumed does not exceed the suggested serving size.
- The benefits of DHM are most effective when they are supported by a healthy diet and plenty of exercise.



812 Meadow Lark Lane, Godlettsville, TN 37072 Telephone: 615-239-8604

Certificate of Analysis

Synaptent LLC 47 W Polk Street, 100-241 Chicago, IL 60654

Product Name	Dihyromyricetin	Client Lot Number	190604
Report Date	02/10/20	Laboratory Number	13625

Test	Method	Result
Identification	H-NMR	Conforms
Assay	HPLC	100.6%
Lead	ICP-MS USP <730>	0.101 ppm
Arsenic	ICP-MS USP <730>	1.364 ppm
Cadmium	ICP-MS USP <730>	<0.0001 ppm
Mercury	ICP-MS USP <730>	0.005 ppm
Total Aerobic count	Biolumix	<1, 000 cfu/g
Yeast and Mold	Biolumix	<100 cfu/g
E. Coli	Biolumix	Negative
Coliform	Biolumix	<10 cfu/g
Salmonella	Biolumix	Negative

Collin Thomas CMOC
Laboratory Manager

02/10/2020 2/10/20 Date

The result(s) stated in this report is only for the sample submitted. This report may not be reproduced in whole or in part, nor may any reference be made to the work, the result, or the company in any news release, public announcements or advertising without our prior written consent.

